

RECONFIRMING THE PRENUCLEOLUS

GUNI ORSHAN AND PETER SUDHÖLTER

By means of an example, it is shown that the prenucleolus is not the only minimal solution that satisfies nonemptiness, Pareto optimality, covariance, the equal treatment property, and the reduced game property, even if the universe of players is infinite. This example also disproves a conjecture of Gurvich et al. (1994). As a second result, we prove that the prenucleolus is axiomatized by nonemptiness, covariance, the equal treatment property, and the reconfirmation property, provided the universe of players is infinite.

1. Introduction. The prenucleolus and the prekernel are widely accepted solutions for cooperative transferable utility games. Introduced as auxiliary solutions of the prebargaining set, they became important solutions in their own rights, heavily supported by the fact that they can be justified by simple and intuitive axioms. Both are closely related because they share many properties and because one, the prenucleolus, is a subsolution of the other. Indeed, both solutions are *nonempty* (NE), *Pareto optimal* (PO), *covariant under strategic equivalence* (COV), *anonymous* (AN), and satisfy the *equal treatment property* (ETP), and the *reduced game property* (RGP). One additional property for each of the solutions suffices to characterize it: The prenucleolus is *single-valued* (SIVA) and the prekernel satisfies the *converse reduced game property* (CRGP). In fact, the prenucleolus is axiomatized by means of SIVA, COV, AN, and RGP (see Sobolev 1975), whereas the prekernel is axiomatized by NE, PO, COV, ETP, RGP, and CRGP (see Peleg 1986). AN can be replaced by ETP in Sobolev's axiomatization (see Orshan 1993) and CRGP in Peleg's axiomatization can be replaced by a *maximality principle*: The prekernel is the maximum solution that satisfies the remaining axioms NE, PO, COV, ETP, and RGP.

In this work we investigate the role of SIVA in the foregoing axiomatizations of the prenucleolus, and we provide two results.

The first result may be motivated as follows. In view of the facts that SIVA may be considered as a “minimality principle” and that the prenucleolus of a game is a distinguished special point of the prekernel, the following question arises in a natural way:

(1) Is it possible to replace SIVA by NE and a *minimality principle*, i.e., is the prenucleolus the *minimum* (or at least the *unique minimal*) solution that satisfies NE, PO, COV, ETP, and RGP?

The first result, Theorem 3.7, shows that the answer to question (1) is negative. Indeed, we construct a solution $\hat{\sigma}$ that is **finite-valued** and satisfies NE, PO, COV, AN, ETP, RGP and that does not contain the prenucleolus. Thus, $\hat{\sigma}$ contains a minimal subsolution having the desired properties (see Corollary 3.8). This result yields a new aspect of the impact of SIVA in Sobolev's (1975) and in Orshan's (1993) axiomatization of the prenucleolus. Hence it “reconfirms” the prenucleolus in the sense that it implicitly reveals a part of the special character of this solution.

As a byproduct, the nonuniqueness result, Theorem 3.7, disproves the following conjecture raised by Gurvich et al. (1994): Any TU game is the reduced game of a “huge” game,

Received September 12, 2001; revised March 18, 2002, and September 26, 2002.

MSC 2000 subject classification. Primary: 91A12.

ORMS subject classification. Primary: Games/group decisions, cooperative.

Key words. Prenucleolus, game theory.

the prekernel of which consists of the prenucleolus only, with respect to the prenucleolus of the “huge” game. If this conjecture were true, then a proof of it would also yield a new proof of Sobolev’s or Orshan’s result, because any solution with the desired properties is a subsolution of the prekernel.

Hence, the prenucleolus cannot be characterized by replacing the “maximality principle” CRGP in the aforementioned axiomatization of the prekernel by a suitable minimality principle. Nevertheless, one may ask the following question:

(2) Is it possible to find an intuitive axiom that characterizes the prenucleolus, if the “maximality principle” CRGP is replaced by this axiom, i.e., is the prenucleolus characterized by NE, PO, COV, ETP, RGP, and some additional intuitive axiom?

The second result, Theorem 4.3, shows that there is a suitable axiom that yields the desired characterization. This axiom requires from any member of the solution of every game that every element of the solution of the reduced game with respect to every coalition, when combined with the restriction of the initial element to the complement coalition, establishes a member of the solution of the initial game. In Balinski and Young (1982) this axiom is one condition of a property they called “uniformity”, and Shimomura (1992) used the term “flexibility”. Recently, Hwang and Sudhölter (2001) called this intuitive axiom the *reconfirmation property* (RCP), which reflects a natural interpretation of the property, and they employed RCP in an axiomatization of the core, which emphasizes the importance of this axiom. Due to private communication, the authors know since 1993 that the prenucleolus is characterized by NE, COV, ETP, RGP, and RCP. Note that PO can be deduced from the first four axioms. Surprisingly, it turns out that RGP is **not** needed in this characterization; that is, the prenucleolus is axiomatized by NE, COV, ETP, and RCP.

We think this axiomatization gives a new insight into the character of the prenucleolus and that it reinforces RCP as a significant property that plays an important role in some characterizations of “classical” solutions like the core and the prenucleolus. Of course, this result also demonstrates the “power” of RCP when combined with ETP. Indeed, AN, even together with PO and RGP, does not replace ETP in Theorem 4.3 because, for example, the *positive* core (see Definition 3.1) satisfies NE, PO, AN, COV, RGP, and RCP. Also it should be remarked that, for single-valued solutions, RGP and RCP are equivalent. If single-valuedness is not required, then there are important solutions that satisfy RCP but violate RGP. For example, the least core satisfies NE, PO, AN, COV, and RCP, but it violates RGP. Particularly this fact demonstrates that RCP stands for itself as an interesting property, independently of RGP.

Though the main theorems, the nonuniqueness result, and the new axiomatization of the prenucleolus are mathematically independent, the above discussion shows their close relationship.

We now briefly review the contents of the paper. Section 2 recalls definitions of some relevant solutions and contains a list of some properties of solutions. In §3, Theorem 3.7 is proved, and §4 contains the new characterization of the prenucleolus and examples that show that each of the employed axioms is logically independent of the remaining axioms.

2. Notation, solutions, and properties. In this section some relevant definitions and results from Maschler et al. (1972) and Peleg (1986) are recalled. Let U be a universe of players containing, without loss of generality, $1, \dots, k$ whenever $|U| \geq k$. A (cooperative TU) *game* is a pair (N, v) such that $\emptyset \neq N \subseteq U$ is finite and $v: 2^N \rightarrow \mathbb{R}$, $v(\emptyset) = 0$. For any game (N, v) let

$$X^*(N, v) = \{x \in \mathbb{R}^N \mid x(N) \leq v(N)\} \text{ and } X(N, v) = \{x \in \mathbb{R}^N \mid x(N) = v(N)\}$$

denote the set of *feasible* and *Pareto optimal* feasible payoffs (*preimputations*), respectively. We use $x(S) = \sum_{i \in S} x_i$ ($x(\emptyset) = 0$) for every $S \in 2^N$ and every $x \in \mathbb{R}^N$ as a convention.

Additionally, x_S denotes the restriction of x to S , i.e., $x_S = (x_i)_{i \in S}$. For disjoint coalitions $S, T \in 2^N$ let $(x_S, x_T) = x_{S \cup T}$. For $x \in \mathbb{R}^N$, $S \subseteq N$, and distinct players $k, l \in N$ let

$$e(S, x, v) = v(S) - x(S) \text{ and } s_{kl}(x, v) = \max_{S \subseteq N \setminus \{l\}: k \in S} e(S, x, v)$$

denote the *excess* of S and the *maximal surplus* of k over l , respectively, at x with respect to (N, v) . The *prekernel* of (N, v) is given by

$$\mathcal{PK}(N, v) = \{x \in X(N, v) \mid s_{kl}(x, v) = s_{lk}(x, v) \ \forall k \in N, \ l \in N \setminus \{k\}\}.$$

For $X \subseteq \mathbb{R}^N$ let $\mathcal{N}((N, v); X)$ denote the *nucleolus* of (N, v) with respect to X , i.e. the set of members of X that lexicographically minimize the nonincreasingly ordered vector of excesses of the coalitions (see Schmeidler 1969). It is well known that the nucleolus with respect to $X^*(N, v)$ is a singleton, the unique element of which is called the *prenucleolus* of (N, v) and is denoted by $\nu(N, v)$.

In general, a *solution* σ associates with each game (N, v) a subset of $X^*(N, v)$. Let σ be a solution. Then σ

(1) is *covariant under strategic equivalence* (COV), if for all games $(N, v), (N, w)$ satisfying $w = \beta v + z$ for some $\beta > 0, z \in \mathbb{R}^N$ the equation $\sigma(N, w) = \beta \sigma(N, v) + z$ holds. (Here we use the convention that identifies $z \in \mathbb{R}^N$ with the *additive* coalitional function, again denoted by z , on the player set N defined by $z(S) = \sum_{i \in S} z_i$ for all $S \in 2^N$. Also note that the games v and w are called *strategically equivalent*.);

(2) is *nonempty* (NE), if $\sigma(N, v) \neq \emptyset$ for every game (N, v) ;

(3) is *Pareto optimal* (PO), if $\sigma(N, v) \subseteq X(N, v)$ for every game (N, v) ;

(4) is *single-valued* (SIVA), if $|\sigma(N, v)| = 1$ for every game (N, v) ;

(5) is *anonymous* (AN), if the following condition is satisfied for all games (N, v) and (M, w) . If $\pi: N \rightarrow M$ is a bijection such that $\pi v = w$, then $\sigma(M, w) = \pi(\sigma(N, v))$. (In this case the games (N, v) and (M, w) are *isomorphic*.);

(6) satisfies the *equal treatment property* (ETP), if for every game (N, v) , for every $x \in \sigma(N, v)$, $x_k = x_l$ for all *substitutes* $k, l \in N$. (Here k and l are *substitutes*, if $v(S \cup \{k\}) = v(S \cup \{l\}) \ \forall S \subseteq N \setminus \{k, l\}$.);

(7) satisfies the *reduced game property* (RGP), if for every game (N, v) , $\emptyset \neq S \subseteq N$, and every $x \in \sigma(N, v)$, $x_S \in \sigma(S, v^{S,x})$. (The *reduced game* $(S, v^{S,x})$ is defined by $v^{S,x}(\emptyset) = 0$, $v^{S,x}(S) = v(N) - x(N \setminus S)$, and $v^{S,x}(T) = \max_{Q \subseteq N \setminus S} (v(T \cup Q) - x(Q))$ for $\emptyset \neq T \subsetneq S$);

(8) satisfies the *converse reduced game property* (CRGP), if for every game (N, v) with $|N| \geq 2$ the following condition is satisfied for every $x \in X(N, v)$: If, for every $S \subseteq N$ with $|S| = 2$, $x_S \in \sigma(S, v^{S,x})$, then $x \in \sigma(N, v)$;

(9) satisfies the *reconfirmation property* (RCP), if for every game (N, v) , $\emptyset \neq S \subseteq N$, for every $x \in \sigma(N, v)$ and $y \in \sigma(S, v^{S,x})$, $(y, x_{N \setminus S}) \in \sigma(N, v)$.

For interpretations and discussions, in particular of the variants 7, 8, and 9 of the reduced game property, see Peleg (1986) and Sudhölter (2001).

We conclude this section by stating the known characterizations of the prenucleolus. Theorem 2.2 will be used in §4.

THEOREM 2.1 (SOBOLEV 1975). *The unique solution that satisfies SIVA, COV, AN, and RGP is the prenucleolus, provided $|U| = \infty$.*

THEOREM 2.2 (ORSHAN 1993). *The unique solution that satisfies SIVA, COV, ETP, and RGP is the prenucleolus, provided $|U| = \infty$.*

3. Nonuniqueness of a minimal solution. This section is devoted to the construction of a solution $\hat{\sigma}$ that satisfies NE, COV, PO, ETP, RGP and that does not contain the prenucleolus as a subsolution. Moreover, as it is finite valued, the solution contains a minimal subsolution satisfying the axioms. As a byproduct, the constructive proof provides an example that disproves the conjecture of Gurvich et al. (1994) mentioned in §1. Throughout this section we shall assume that $|U| \geq 4$.

First, a specific *coalition structure*, i.e., a partition of the set of players, is defined. Let (N, v) be a game and $x = \nu(N, v)$. Define the binary relation \sim_v on N by

$$k \sim_v l \Leftrightarrow k = l \text{ or } (k \neq l \text{ and } s_{kl}(x, v) \leq 0)$$

and note that \sim_v is reflexive and transitive. It is also symmetric (an equivalence relation), because $x \in \mathcal{PH}(N, v)$. Let $\mathcal{T}(N, v)$ denote the set of equivalence classes of \sim_v . In Section 3 of Maschler et al. (1972), the collection $\mathcal{T}(N, v)$ of coalitions is the partition, which corresponds to the smallest excess strictly greater than 0, of the *profile* generated by the prekernel element $x = \nu(N, v)$. (Of course, $\mathcal{T}(N, v) = \{N\}$ in the case that there is no coalition of positive excess.)

DEFINITION 3.1. Let (N, v) be a game and $x = \nu(N, v)$. Let $t^+ = \max\{t, 0\}$ denote the positive part of the real number t . The *positive core* of (N, v) is the set

$$\mathcal{C}^+(N, v) = \{y \in X^*(N, v) \mid (e(S, x, v))^+ = (e(S, y, v))^+ \ \forall S \subseteq N\}.$$

Note that the positive core was first mentioned by Orshan (1994), who fully described this solution in the three-person case. Though this solution may be regarded as an interesting core extension, in the present paper it serves as an auxiliary solution only.

LEMMA 3.2. *The positive core satisfies RGP and RCP.*

Though this lemma is contained in Sudhölter (1993, Lemma 2.2), we shall present a proof that proceeds analogously to the proof of RGP of the prenucleolus due to Peleg (1988).

PROOF. Let (N, v) be a game, $x \in X(N, v)$ and $\emptyset \neq S \subseteq N$. For $\alpha \in \mathbb{R}$ define

$$\mathcal{D}(\alpha, x, v) = \{S \subseteq N \mid e(S, x, v) \geq \alpha\} \cup \{\emptyset, N\}.$$

Then Definition 3.1 can be formulated as

$$(3.1) \quad x \in \mathcal{C}^+(N, v) \Leftrightarrow \mathcal{D}(\alpha, x, v) = \mathcal{D}(\alpha, \nu(N, v), v) \ \forall \alpha > 0.$$

According to Kohlberg (1971), the prenucleolus is characterized by the equivalence

$$(3.2) \quad x = \nu(N, v) \Leftrightarrow \left(\forall \alpha \in \mathbb{R} \ \forall y \in \mathbb{R}^N : \begin{array}{l} y(N) = 0, \ y(T) \geq 0 \ \forall T \in \mathcal{D}(\alpha, x, v) \\ \Rightarrow y(T) = 0 \ \forall T \in \mathcal{D}(\alpha, x, v) \end{array} \right).$$

By Equations (3.1) and (3.2), the positive core is characterized by the equivalence

$$(3.3) \quad x \in \mathcal{C}^+(N, v) \Leftrightarrow \left(\forall \alpha > 0 \ \forall y \in \mathbb{R}^N : \begin{array}{l} y(N) = 0, \ y(T) \geq 0 \ \forall T \in \mathcal{D}(\alpha, x, v) \\ \Rightarrow y(T) = 0 \ \forall T \in \mathcal{D}(\alpha, x, v) \end{array} \right).$$

Note that $x_S \in X(N, v^{S,x})$ and

$$(3.4) \quad \mathcal{D}(\alpha, x_S, v^{S,x}) = \{S \cap T \mid T \in \mathcal{D}(\alpha, x, v)\} \ \forall \alpha \in \mathbb{R}.$$

By Equation (3.4), for any $\alpha \in \mathbb{R}$, the set

$$\{y_S \in \mathbb{R}^S \mid y_S(S) = 0, \ y_S(Q) \geq 0 \ \forall Q \in \mathcal{D}(\alpha, x_S, v^{S,x})\}$$

is the projection of

$$\{y \in \mathbb{R}^N \mid y(N) = 0, y_i = 0 \forall i \in N \setminus S, y(T) \geq 0 \forall T \in \mathcal{D}(\alpha, x, v)\}.$$

Thus RGP is implied by Equation (3.3). To show RCP let $x \in \mathcal{C}^+(N, v)$ and $z \in \mathcal{C}^+(S, v^{S,x})$. By RGP, $x_S \in \mathcal{C}^+(S, v^{S,x})$, thus $\mathcal{D}(\alpha, x_S, v^{S,x}) = \mathcal{D}(\alpha, z, v^{S,x}) \forall \alpha > 0$ by (3.1). Hence, by Equation (3.4) applied to $(z, x_{N \setminus S})$, $\mathcal{D}(\alpha, (z, x_{N \setminus S}), v) = \mathcal{D}(\alpha, x, v)$ for every $\alpha > 0$. Thus Equation (3.3) finishes the proof. \square

REMARK 3.3. Let (N, v) be a game and $x = \nu(N, v)$. By the definition of the positive core we obtain

$$\forall k, l \in N, k \neq l: \left(\begin{array}{l} s_{kl}(x, v) > 0 \Rightarrow s_{kl}(x, v) = s_{kl}(y, v) \text{ and} \\ s_{kl}(x, v) \leq 0 \Leftrightarrow s_{kl}(y, v) \leq 0 \end{array} \right) \forall y \in \mathcal{C}^+(N, v).$$

Let (N, v) be a game. For any total order \leq of $\mathcal{T} := \mathcal{T}(N, v)$, let us say $\mathcal{T} = \{T_1, \dots, T_t\}$ with $T_1 < \dots < T_t$, recursively define $\alpha_{\leq}^v \in \mathbb{R}^{\mathcal{T}}$ (we abbreviate α_{\leq}^v by α if there is no danger of confusion) by

$$(3.5) \quad \alpha(T_1) = \min \{y(T_1) \mid y \in \mathcal{C}^+(N, v)\}$$

and

$$(3.6) \quad \alpha(T_i) = \min \{y(T_i) \mid y \in \mathcal{C}^+(N, v), y(T_j) = \alpha(T_j) \forall j = 1, \dots, i-1\}$$

for all $i = 2, \dots, t$ and put

$$(3.7) \quad X_{\leq}(N, v) = \{z \in \mathbb{R}^N \mid z(T) = \alpha_{\leq}^v(T) \forall T \in \mathcal{T}(N, v)\}.$$

REMARK 3.4. Let (N, v) be a game. Note that α_{\leq}^v is well defined because $\mathcal{C}^+(N, v)$ is nonempty and compact. Therefore $X_{\leq}(N, v)$ is a nonempty convex set of preimputations, which, by Equations (3.5), (3.6), and (3.7), intersects the positive core; i.e., we have

$$(3.8) \quad \mathcal{C}_{\leq}^+(N, v) := \mathcal{C}^+(N, v) \cap X_{\leq}(N, v) \neq \emptyset.$$

This subset of the positive core can also be expressed as

$$(3.9) \quad \mathcal{C}_{\leq}^+(N, v) = \{y \in \mathcal{C}^+(N, v) \mid (y(T_1), \dots, y(T_t)) \leq_{lex} (z(T_1), \dots, z(T_t)) \forall z \in \mathcal{C}^+(N, v)\}.$$

LEMMA 3.5. Let (N, v) be a game and \leq be a total order of $\mathcal{T}(N, v)$. Then the nucleolus of (N, v) with respect to $X_{\leq}(N, v)$ is a singleton, which belongs to $\mathcal{C}^+(N, v)$.

PROOF. By Equation (3.8), $\mathcal{C}_{\leq}^+(N, v) \neq \emptyset$. Hence,

$$(3.10) \quad \mathcal{N}((N, v); X_{\leq}(N, v)) = \mathcal{N}((N, v); \mathcal{C}_{\leq}^+(N, v))$$

by the definition of the positive core. By Equation (3.9) the latter set of preimputations is a nonempty, convex, and compact set. Thus $|\mathcal{N}((N, v); \mathcal{C}_{\leq}^+(N, v))| = 1$ (see Schmeidler 1969). \square

DEFINITION 3.6. For any game (N, v) and any total order \leq of $\mathcal{T}(N, v)$ let $\nu_{\leq}(N, v)$ be the unique member of $\mathcal{N}((N, v); X_{\leq}(N, v))$. The solution $\widehat{\sigma}$ is defined by

$$\widehat{\sigma}(N, v) = \{\nu_{\leq}(N, v) \mid \leq \text{ is a total order of } \mathcal{T}(N, v)\}.$$

THEOREM 3.7. The solution $\widehat{\sigma}$ satisfies NE, PO, COV, ETP, RGP, and AN. Moreover, the prenucleolus is not a subsolution of $\widehat{\sigma}$.

PROOF. Let (N, v) be a game and \preceq be a total order of $\mathcal{T} = \mathcal{T}(N, v)$.

(1) We show that $\widehat{\sigma}$ satisfies NE, AN, PO, and COV. The vector $x := \nu_{\preceq}(N, v)$ is a member of $\widehat{\sigma}(N, v)$, thus $\widehat{\sigma}$ satisfies NE. A bijection $\pi: N \rightarrow M$ maps \mathcal{T} bijectively to $\mathcal{T}(M, \pi v)$, because the prenucleolus satisfies AN. Thus $\widehat{\sigma}$ satisfies AN as well. Moreover, $\widehat{\sigma}$ satisfies PO, because it is a subsolution of \mathcal{C}^+ . Let $\beta > 0$, $z \in \mathbb{R}^N$, $w := \beta v + z$, and $y := \beta x + z$. Then $\mathcal{T}(N, w) = \mathcal{T}$, because the prenucleolus satisfies COV. Moreover, by COV of the positive core (which can be shown in a straightforward way), the equation

$$\mathcal{C}_{\preceq}^+(N, w) = \beta \mathcal{C}_{\preceq}^+(N, v) + z$$

is valid, thus $\nu_{\preceq}(N, w) = \beta \nu_{\preceq}(N, v) + z = y$ by Equation (3.10). Hence $\widehat{\sigma}$ satisfies COV.

(2) To show ETP and RGP a *derived* game (N, v_{\preceq}) is defined by

$$v_{\preceq}(S) = \begin{cases} v(S), & \text{if } S \in 2^N \setminus \mathcal{T}, \\ \alpha_{\preceq}^v(T), & \text{if } S = T \text{ for some } T \in \mathcal{T}. \end{cases}$$

Then

$$(3.11) \quad \mathcal{N}((N, v); X_{\preceq}(N, v)) = \mathcal{N}((N, v_{\preceq}); X_{\preceq}(N, v)),$$

because v_{\preceq} differs from v at most on the partition \mathcal{T} , the elements T of which receive a fixed amount $z(T) = \alpha_{\preceq}^v(T)$ by every preimputation z of $X_{\preceq}(N, v)$. The expression on the right-hand side of Equation (3.11) is the prenucleolus $\nu((N, v_{\preceq}); \mathcal{T})$ of the game (N, v_{\preceq}) with the *coalition structure* \mathcal{T} . Thus,

$$(3.12) \quad x := \nu_{\preceq}(N, v) = \nu((N, v_{\preceq}); \mathcal{T}).$$

It is well known that the prenucleolus of a game with coalition structure is a member of the prekernel of the game with coalition structure. For every pair (k, l) of distinct players the equation

$$(3.13) \quad s_{kl}(x, v) = \begin{cases} s_{kl}(x, v_{\preceq}), & \text{if } k, l \in T \text{ for some } T \in \mathcal{T}, \\ s_{kl}(\nu(N, v), v), & \text{otherwise,} \end{cases}$$

is a consequence of the definition of the derived game and of Lemma 3.2 and Remark 3.3, respectively. Equations (3.12) and (3.13) imply $x \in \mathcal{PK}(N, v)$. Thus $\widehat{\sigma}$ is a subsolution of the prekernel which satisfies ETP.

To show that $\widehat{\sigma}$ satisfies RGP, let $\emptyset \neq S \subseteq N$ and $w = v^{S,x}$. Using the well-known fact

$$s_{kl}(x_S, w) = s_{kl}(x, v) \quad \forall k, l \in S \text{ with } k \neq l,$$

we obtain

$$(3.14) \quad \mathcal{T}(S, w) = \{T \cap S \mid T \in \mathcal{T} \text{ and } T \cap S \neq \emptyset\}.$$

Let \preceq^S be the total order on $\mathcal{T}(S, w)$ consistent with \preceq , which is defined by the following requirement: If $T, T' \in \mathcal{T}$ satisfy $T < T'$ and $T \cap S \neq \emptyset \neq T' \cap S$, then $T \cap S <^S T' \cap S$. By RGP and RCP of the positive core (see Lemma 3.2),

$$\mathcal{C}_{\preceq^S}^+(S, w) = \{y \in \mathbb{R}^S \mid (y, x_{N \setminus S}) \in \mathcal{C}_{\preceq}^+(N, v)\},$$

and thus,

$$X_{\preceq^S}(S, w) = \{y \in \mathbb{R}^S \mid (y, x_{N \setminus S}) \in X_{\preceq}(N, v)\}.$$

Therefore, $(v_{\preceq})^{S,x} = w_{\preceq^S}$. Hence $\nu_{\preceq^S}(S, w) = x_S$ by RGP of the prenucleolus of games with coalition structures (see Peleg (1988, Theorem 5.2.7)).

(3) To show that ν is not a subsolution of $\widehat{\sigma}$ the following “cyclic” four-person game (M, u) is defined by $M = \{1, \dots, 4\}$ and

$$u(S) = \begin{cases} 1, & \text{if } S \in \{\{1, 2\}, \{2, 3\}, \{3, 4\}, \{4, 1\}\}, \\ 0, & \text{if } S \in \{\emptyset, M\}, \\ -2, & \text{otherwise.} \end{cases}$$

Note that (M, u) is *transitive*. (A game is transitive if its symmetry group—i.e., the group of permutations of N that do not change the game—is transitive.) Indeed, the cyclic permutation, which maps 1 to 2, 2 to 3, 3 to 4, and 4 to 1, is a symmetry. Hence, by AN and PO, we obtain $\nu(M, u) = 0 \in \mathbb{R}^M$. Therefore, $x \in \mathcal{C}^+(M, u)$, iff $x(S) = 0$ for every $S = \{1, 2\}, \dots, \{4, 1\}$, and $x(T) \geq -2$ for every $T \subseteq M$. These inequalities show that $\mathcal{C}^+(M, u) = \text{convex hull}\{(-1, 1, -1, 1), (1, -1, 1, -1)\}$. Also, $\mathcal{F}(M, u) = \{\{k\} \mid k \in M\}$. Thus $\widehat{\sigma}(M, u) = \{(-1, 1, -1, 1), (1, -1, 1, -1)\}$. \square

COROLLARY 3.8. *There is a minimal solution that satisfies NE, PO, COV, ETP, RGP, AN, and that does not coincide with the prenucleolus.*

PROOF. Let Σ denote the partially ordered set of subsolutions of $\widehat{\sigma}$ that satisfy NE, PO, COV, ETP, RGP, and AN. By Theorem 3.7, $\widehat{\sigma} \in \Sigma$. To show that a chain Σ^0 (a subset of comparable elements) has a lower bound we verify that σ^0 , defined by $\sigma^0(N, v) = \bigcap_{\sigma \in \Sigma^0} \sigma(N, v)$, belongs to Σ . The solution σ^0 satisfies PO, COV, ETP, RGP, and AN, because all members of the chain satisfy these axioms. Moreover, σ^0 satisfies NE, because any $\sigma \in \Sigma$ is finite-valued. Hence, by Zorn’s lemma, Σ has a minimal element. By Theorem 3.7, $\nu \notin \Sigma$. \square

In Gurvich et al. (1994, §4.4) the following question is raised: Let (N, v) be a game. Is there any game $(\widetilde{N}, \widetilde{v})$ such that (a) $N \subseteq \widetilde{N}$, (b) $\mathcal{PH}(\widetilde{N}, \widetilde{v})$ is a singleton consisting of the prenucleolus y only, and (c) $\widetilde{v}^{N,y} = v$? A positive answer to this question would yield a new proof Sobolev’s or Orshan’s axiomatization of the prenucleolus. Theorem 3.7 shows that the answer to this question is negative. Moreover, the game (M, u) defined in the last part of the proof of this theorem is an explicit “counter” example. (Note that a one-parameter set of games that contains (M, u) is discussed in Orshan 1994 and a variant of (M, u) is used to prove the main result of Sughölter and Peleg 2002.)

4. An axiomatization of the prenucleolus. This section is devoted to show that the prenucleolus is axiomatized by NE, COV, ETP, and RCP, provided $|U| = \infty$.

Let σ be a solution. Let Γ^2 be the set of games with at most two persons. The following lemmata are useful.

LEMMA 4.1. *Assume that $|U| \geq 2$ and that σ satisfies NE, COV, ETP, and RCP. Then $\sigma(N, v) = \nu(N, v)$ for every $(N, v) \in \Gamma^2$.*

PROOF. Let (M, u) be a two-person game. By NE there exists $x \in \sigma(M, u)$. By ETP and COV there exists $a \in \mathbb{R}$ such that $x_i = u(\{i\}) + a$ for $i \in M$. Let $i \in M$ and $u^i = u^{\{i\},x}$. By NE there exists $y \in \sigma(\{i\}, u^i)$. By COV, $y^\alpha := \alpha(y - u^i(\{i\})) + u^i(\{i\}) \in \sigma(\{i\}, u^i)$ for every $\alpha > 0$. By RCP, $(y^\alpha, x_{M \setminus \{i\}}) \in \sigma(M, u)$, thus $y^\alpha = x_i$ for all $\alpha > 0$. We conclude that $y^\alpha = y$ for all $\alpha > 0$, hence the proof is complete. \square

For any game (N, v) and any $x \in \mathbb{R}^N$ let $\mu(x, v) = \max_{S \subseteq N} e(S, x, v)$.

LEMMA 4.2. *Under the assumptions of Lemma 4.1 and the additional assumption that $|U| = \infty$, the following assertions are valid:*

(1) σ satisfies PO.

(2) *Let (N, v) be a game, let $x \in \sigma(N, v)$, and let $i \in N$. Then there exist $S^i, S^{-i} \subseteq N$ such that $i \in S^i$, $i \notin S^{-i}$, and $e(S^i, x, v) = e(S^{-i}, x, v) = \mu(x, v)$.*

PROOF. Let (N, v) be a game. As $|U| = \infty$ we may assume that $N \subseteq U \setminus \{1, 2\}$ and that $\tilde{N} = \{1, 2\} \cup N \subseteq U$. By NE there exists $x \in \sigma(N, v)$. Define $\tilde{v}(S)$, $S \subseteq \tilde{N}$, by $\tilde{v}(S) = v(S \setminus \{1, 2\})$. (That is, (\tilde{N}, \tilde{v}) arises from (N, v) by adding the two null-players 1 and 2.) By NE there exists $\tilde{x} \in \sigma(\tilde{N}, \tilde{v})$. By ETP, $\tilde{x}_1 = \tilde{x}_2$.

CLAIM 1. $\tilde{x}(\tilde{N}) = \tilde{v}(\tilde{N})$: Let $\tilde{v}^1 = \tilde{v}^{(1), \tilde{x}}$. By Lemma 4.1, $\sigma(\{1\}, \tilde{v}^1) = \{\tilde{v}^1(\{1\})\}$. By RCP, $(\tilde{v}^1(\{1\}), \tilde{x}_{N \setminus \{1\}}) \in \sigma(\tilde{N}, \tilde{v})$. By ETP, $\tilde{v}^1(\{1\}) = \tilde{x}_2 = \tilde{x}_1$, thus $\tilde{x}(\tilde{N}) = \tilde{v}(\tilde{N})$.

CLAIM 2. $\tilde{x}_1 = \tilde{x}_2 = 0$: For every $i \in N$ let $\tilde{v}^{i,1} = \tilde{v}^{(1,i)}$. Assume the contrary. Then two cases may occur:

(1) $\tilde{x}_1 = \tilde{x}_2 < 0$: Then $e(\{1\}, \tilde{x}, \tilde{v}) = -\tilde{x}_1 > 0$, thus $\mu(\tilde{x}, \tilde{v}) > 0$. Let $S \subseteq \tilde{N}$ attain $\mu(\tilde{x}, \tilde{v})$. By Claim 1, $\emptyset \neq S \neq \tilde{N}$. By our assumption, $\{1, 2\} \subseteq S$. Let $i \in N \setminus S$. By Lemma 4.1, $\tilde{y} := \nu(\{1, i\}, \tilde{v}^{i,1}) \in \sigma(\{1, i\}, \tilde{v}^{i,1})$. Then $\tilde{y}_1 > \tilde{x}_1$. By RCP we have $(\tilde{y}, \tilde{x}_{\tilde{N} \setminus \{1, i\}}) \in \sigma(\tilde{N}, \tilde{v})$. A contradiction to ETP is obtained, because $\tilde{y}_1 > \tilde{x}_1 = \tilde{x}_2$.

(2) $\tilde{x}_1 = \tilde{x}_2 > 0$: Then $e(\tilde{N} \setminus \{1\}, \tilde{x}, \tilde{v}) = \tilde{x}_1 > 0$ by Claim 1, thus $\mu(\tilde{x}, \tilde{v}) > 0$. Let $S \subseteq \tilde{N}$ attain $\mu(\tilde{x}, \tilde{v})$. By Claim 1, $\emptyset \neq S \neq \tilde{N}$. By our assumption, $\{1, 2\} \cap S = \emptyset$. Let $i \in S$. By Lemma 4.1, $\tilde{y} := \nu(\{1, i\}, \tilde{v}^{i,1}) \in \sigma(\{1, i\}, \tilde{v}^{i,1})$. Then $\tilde{y}_1 < \tilde{x}_1$. By RCP, $(\tilde{y}, \tilde{x}_{\tilde{N} \setminus \{1, i\}}) \in \sigma(\tilde{N}, \tilde{v})$. A contradiction to ETP is obtained, because $\tilde{y}_1 < \tilde{x}_1 = \tilde{x}_2$.

Now assertion (1) of our lemma can be deduced. By Claim 2, $(N, \tilde{v}^{N, \tilde{x}}) = (N, v)$, thus $\tilde{x} := (0, 0, x) \in \sigma(\tilde{N}, \tilde{v})$ by RCP. By Claim 1, $x(N) = v(N)$.

To prove assertion (2), we have to show that the following conditions are satisfied:

$$(4.1) \quad N = \bigcup \{S \subseteq N \mid e(S, x, v) = \mu(x, v)\},$$

$$(4.2) \quad \emptyset = \bigcap \{S \subseteq N \mid e(S, x, v) = \mu(x, v)\}.$$

Assume the contrary. Then two cases may occur:

(1) There exists $i \in N \setminus \bigcup \{S \subseteq N \mid e(S, x, v) = \mu(x, v)\}$: By Lemma 4.1,

$$\tilde{y} = \nu(\{1, i\}, \tilde{v}^{(1,i), \tilde{x}}) \in \sigma(\{1, i\}, \tilde{v}^{(1,i), \tilde{x}}).$$

The fact that $\tilde{y}_1 > 0 = \tilde{x}_2$, is in contradiction to ETP.

(2) There exists $i \in \bigcap \{S \subseteq N \mid e(S, x, v) = \mu(x, v)\}$: By Lemma 4.1,

$$\tilde{y} = \nu(\{1, i\}, \tilde{v}^{(1,i), \tilde{x}}) \in \sigma(\{1, i\}, \tilde{v}^{(1,i), \tilde{x}}).$$

The fact that $\tilde{y}_1 < 0 = \tilde{x}_2$, is in contradiction to ETP. \square

Now the main theorem of this section can be proved.

THEOREM 4.3. *The prenucleolus is the unique solution that satisfies NE, COV, ETP, and RCP, provided $|U| = \infty$.*

PROOF. The prenucleolus satisfies the desired properties. Indeed, it satisfies RCP, because RCP and RGP are equivalent for single-valued solutions. To show the opposite direction let σ be a solution that satisfies the desired axioms. By Lemma 4.1, σ satisfies PO.

In view of Theorem 2.2 it suffices to show that σ satisfies SIVA. Let (N, v) be a game. Take a disjoint copy $N^* \subseteq U$ of N , i.e.,

$$N \cap N^* = \emptyset \text{ and } N \rightarrow N^*, i \mapsto i^*, \text{ is a bijection.}$$

Choose any real number α satisfying $\alpha > (n^2 + n) \max_{P, Q \subseteq N} (v(P) - v(Q))$ and define a “replicated” game $(N \cup N^*, \hat{v})$ by

$$\hat{v}(S \cup T^*) = \begin{cases} v(S), & \text{if } T = S, \\ -\alpha, & \text{otherwise,} \end{cases}$$

where $S, T \subseteq N$. Let $z \in \sigma(N \cup N^*, \hat{v})$. It is the aim to show that the reduced game (N, u) with respect to N and z (defined by $u = \hat{v}^{N, z}$) is given by

$$(4.3) \quad u(S) = \hat{v}(S \cup S^*) - z(S) = v(S) - z(S) \quad \forall S \subseteq N.$$

To prove (4.3), first note that for any $i \in N$ the players i and i^* are substitutes. Hence, by ETP, $z_i = z_{i^*}$ for all $i \in N$.

CLAIM 1. For all $i \in N$, $z_i \geq \min_{P, Q \subseteq N} (v(P) - v(Q))$:

Assume, on the contrary, that there exists $i_0 \in N$ such that $z_{i_0} < \min_{P, Q \subseteq N} (v(P) - v(Q))$. Choose any coalition $S \cup T^*$ attaining $\mu(z, \hat{v})$. In view of the fact that

$$e(\{i_0, i_0^*\}, z, \hat{v}) = v(\{i_0\}) - 2z_{i_0} > -z_{i_0} > 0,$$

the maximal excess cannot be attained by \emptyset or by $N \cup N^*$. By Lemma 4.2, Claim 1 is shown as soon as $i_0 \in S$ is verified.

Assume, on the contrary, $i_0 \notin S$. If $i_0 \notin T$, then the observation

$$e(S \cup \{i_0\} \cup T^* \cup \{i_0^*\}, z, \hat{v}) - e(S \cup T^*, z, \hat{v}) = \begin{cases} -2z_{i_0} > 0, & \text{if } S \neq T, \\ v(S \cup \{i_0\}) - v(S) - 2z_{i_0} > 0, & \text{if } S = T \end{cases}$$

yields the desired contradiction in this case. If $i_0 \in T$, then the observation

$$e(S \cup \{i_0\} \cup T^*, z, \hat{v}) - e(S \cup T^*, z, \hat{v}) = \begin{cases} -z_{i_0} > 0, & \text{if } S \neq T \setminus \{i_0\}, \\ v(S \cup \{i_0\}) + \alpha - z_{i_0} > 0, & \text{if } S = T \setminus \{i_0\} \end{cases}$$

yields the desired contradiction.

CLAIM 2. $z_i \leq n \max_{P, Q \subseteq N} (v(P) - v(Q)) \quad \forall i \in N$:

Let $i_0 \in N$ be a player. Observe that

$$v(N) = 2z(N) = 2z_{i_0} + 2z(N \setminus \{i_0\}) \geq 2z_{i_0} + 2(n-1) \min_{P, Q \subseteq N} (v(P) - v(Q))$$

by PO, ETP, and Claim 1. Thus our claim follows immediately.

Now the proof can be finished. Put $\tilde{S} = \{i \in N \mid z_i < 0\}$ and observe that

$$(4.4) \quad u(S) = \max\{v(S) - z(S), -\alpha - z(\tilde{S})\} \quad \forall \emptyset \neq S \subsetneq N.$$

Let S be a nontrivial ($\emptyset \neq S \subsetneq N$) coalition. Then

$$v(S) - z(S) \geq v(S) - (n-1) \max_{P, Q \subseteq N} (v(P) - v(Q)) \geq -n \max_{P, Q \subseteq N} (v(P) - v(Q))$$

and

$$-\alpha - z(\tilde{S}) \leq -\alpha + n^2 \max_{P, Q \subseteq N} (v(P) - v(Q)) < -n \max_{P, Q \subseteq N} (v(P) - v(Q)),$$

where the last inequality is implied by the definition of α . Hence u is given by

$$u(S) = v(S) - z(S) \quad \forall S \subseteq N.$$

By NE there exists $x \in \sigma(N, v)$. COV implies $x - z_N \in \sigma(N, u)$, thus $(x - z_N, z_{N^*}) \in \sigma(N \cup N^*, \hat{v})$ by RCP. ETP implies $x - z_N = z_N$, thus $x = 2z_N$ is the unique member of $\sigma(N, v)$. \square

Four examples are presented that show that each of the axioms (1) NE, (2) COV, (3) ETP, and (4) RCP is logically independent of the remaining axioms in Theorem 4.3.

Let (N, v) be a game. Let σ^i , $i = 1, 2$, be defined by

$$\begin{aligned} \sigma^1(N, v) &= \emptyset, \\ \sigma^2(N, v) &= \{x \in X(N, v) \mid x_i = x_j \quad \forall i, j \in N\}. \end{aligned}$$

Let \leq be a total order of U . For every finite set $N \subseteq U$ let \leq_N be the restriction of \leq to N and let \leq_{lex}^N be the induced lexicographical order on \mathbb{R}^N . Then σ^3 is defined by

$$\sigma^3(N, v) = \{x \in \mathcal{C}^+(N, v) \mid x \leq_{lex}^N y \ \forall y \in \mathcal{C}^+(N, v)\}.$$

Also, let σ^4 be the Shapley value.

It is straightforward to check that σ^i , $i = 1, \dots, 4$, satisfies all properties except the i th one. If $|U| \geq 4$, then none of the solutions coincides with the prenucleolus.

The examples also show that each axiom employed in Sobolev's or Orshan's characterization of the prenucleolus is logically independent of the remaining axioms.

REMARK 4.4. The prekernel satisfies NE, COV, AN, ETP, and RGP. Thus RCP cannot be replaced by RGP in Theorem 4.3. The positive core (see Definition 3.1) satisfies NE, COV, AN, RGP, and RCP, hence ETP cannot be replaced by AN (as in the "classical") axiomatization.

Another well-known solution is the least core. The *least core* of a game (N, v) is defined by

$$\mathcal{LC}(N, v) = \left\{ x \in X(N, v) \mid \max_{\emptyset \neq S \subsetneq N} e(S, x, v) = \min_{\emptyset \neq S \subsetneq N} e(S, \nu(N, v), v) \right\}.$$

It is well known and easy to verify that \mathcal{LC} satisfies NE, COV, and AN. It also satisfies RCP because the maximal excess of nontrivial coalitions in a reduced game is not larger than the maximal excess of nontrivial coalitions in the game. The least core does not satisfy RGP.

REMARK 4.5. The infinity assumption on $|U|$ in Theorem 4.3 is crucial. Indeed, if $|U| < \infty$, then define for any game (N, v) ,

$$\sigma(N, v) = \begin{cases} \{\nu(N, v)\}, & \text{if } N \subsetneq U, \\ \{x \in \mathcal{PK}(N, v) \mid x_S = \nu(S, v^{S,x}) \ \forall \emptyset \neq S \subsetneq N\}, & \text{if } N = U. \end{cases}$$

Then σ satisfies all axioms of Theorem 4.3. Also, there are examples that show that the prenucleolus is a proper subsolution of σ when $|U| \geq 4$.

Acknowledgments. The authors are grateful to the referees of this paper for their comments. The second author was supported by the Center for Rationality and Interactive Decision Theory at The Hebrew University of Jerusalem and by the Edmund Landau Center for Research in Mathematical Analysis and Related Areas, sponsored by the Minerva Foundation (Germany).

References

- Balinski, M., H. P. Young. 1982. *Fair Representation*, Yale University Press, New Haven, CT.
- Gurvich, V. A., O. R. Menshikova, I. S. Menshikov. 1994. On the nucleolus, balanced sets of coalitions and Sobolev's theorem. Mimeograph.
- Hwang, Y-A., P. Sudhölter. 2001. Axiomatizations of the core on the universal domain and other natural domains. *Internat. J. Game Theory* **29** 597–623.
- Kohlberg, E. 1971. On the nucleolus of a characteristic function game. *SIAM J. Appl. Math.* **20** 62–66.
- Maschler, M., B. Peleg, L. S. Shapley. 1972. The kernel and bargaining set for convex games. *Internat. J. Game Theory* **1** 73–93.
- Orshan, G. 1993. The prenucleolus and the reduced game property: Equal treatment replaces anonymity. *Internat. J. Game Theory* **22** 241–248.
- . 1994. Non-symmetric prekernels. Discussion Paper 60, Center for Rationality, The Hebrew University of Jerusalem, Jerusalem, Israel.
- Peleg, B. 1986. On the reduced game property and its converse. *Internat. J. Game Theory* **15** 187–200.
- . 1988. Introduction to the theory of cooperative games, Chapter 5. Research Memorandum 84, Center for Research in Mathematical Economics and Game Theory, The Hebrew University of Jerusalem, Jerusalem, Israel.

- Schmeidler, D. 1969. The nucleolus of a characteristic function game. *SIAM J. Appl. Math.* **17** 1163–1170.
- Shimomura, K.-I. 1992. Individual rationality and collective optimality. Ph.D. thesis, University of Rochester, Rochester, New York.
- Sobolev, A. I. 1975. The characterization of optimality principles in cooperative games by functional equations (Russian). N. N. Vorobiev, ed. *Mathematical Methods in the Social Sciences*, Vol. 6, Academy of Sciences of the Lithuanian SSR, Vilnius, Lithuania, 95–151.
- Sudhölter, P. 1993. Independence for characterizing axioms of the pre-nucleolus. Working Paper 220, Institute of Mathematical Economics, University of Bielefeld, Bielefeld, Germany.
- , B. Peleg. 2002. A note on an axiomatization of the core of market games. *Math. Oper. Res.* **27** 441–444.

G. Orshan: Dept. of Agriculture Economics and Management, The Hebrew University of Jerusalem, P.O. Box 12, Rehovot, 76100, Israel, and Dept. of Mathematics, The Open University of Israel, 16 Klausner Street, P.O.B. 39382, Ramat Aviv, Tel Aviv, 61392, Israel; e-mail: orshan@agri.huji.ac.il

P. Sudhölter: Department of Economics, University of Southern Denmark, Campusvej 55, DK-5230 Odense M, Denmark; e-mail: psu@sam.sdu.dk